

## Uses, Side effects, Dosage, Price & Details Of Nimesulide 100mg + Paracetamol 325mg Tablets (YANA-P)

### Uses:

- **Pain Relief:** This combination medication is commonly used to relieve pain, including mild to moderate pain such as headache, toothache, musculoskeletal pain, and menstrual cramps.
- **Fever Reduction:** It is also used to reduce fever associated with various conditions such as infections or inflammatory processes.

### Dosage:

- The dosage of Nimesulide 100mg + Paracetamol 325mg tablets should be determined by a healthcare professional based on the individual's age, weight, medical condition, and response to treatment.
- Generally, the typical dosage for adults is one tablet taken orally every 6 to 8 hours as needed for pain or fever relief.
- The maximum daily dose should not exceed the prescribed limit to avoid the risk of overdose or adverse effects.

### Side Effects:

- Common side effects of this combination medication may include:
  - Nausea and vomiting
  - Upset stomach or abdominal pain
  - Dizziness or lightheadedness
  - Headache
  - Skin rash or itching
- Less common but potentially serious side effects may include:
  - Allergic reactions (rash, itching, swelling, difficulty breathing)
  - Liver damage (especially with prolonged use or overdose of paracetamol)
  - Kidney damage (especially with prolonged use or in individuals with pre-existing kidney problems)
  - Gastrointestinal bleeding or ulcers (especially with prolonged use or in individuals with a history of gastrointestinal disorders)

### Precautions:

- Individuals with certain medical conditions such as liver disease, kidney disease, gastrointestinal disorders, or asthma should use this medication with caution and under the supervision of a healthcare professional.
- Avoid consuming alcohol while taking this medication, as it may increase the risk of liver damage or gastrointestinal side effects.

- Do not exceed the recommended dosage or take this medication for longer than prescribed, as it may increase the risk of adverse effects.
- Inform your healthcare provider about all medications you are taking, including prescription drugs, over-the-counter medications, and herbal supplements, to avoid potential drug interactions.

**Important Note:**

It's crucial to use **Nimesulide 100mg + Paracetamol 325mg tablets** only as prescribed by a healthcare professional. If you experience any concerning side effects or have questions about the medication, consult your healthcare provider promptly.